



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

moon's secular acceleration over and above the late determination. We have seen that the hypothesis of a retardation of the tidal wave by friction of only ten minutes is sufficient, according to Fourier's theory, of the loss of heat, and if we suppose the loss to be somewhat greater than this theory requires, it is only necessary to make the hypothesis a very little greater. If  $6''$  of the moon's secular acceleration is due to tidal action, then the length of the day has increased about  $\frac{1}{140}$  of a second in the last twenty-five hundred years. If the effect of tidal action is insensible, and the late determination of the moon's secular acceleration from theory be received, then we have no way of accounting for the remaining  $6''$ , and besides we cannot allow that the earth's volume has contracted as much as even Fourier's theory requires. The main point at which we have arrived is, that tidal action is adequate to account for  $6''$  of secular acceleration upon a very reasonable and probable hypothesis with regard to the magnitude and retardation of the tidal wave by friction, after making due allowance of the effect for counteracting the effect of a probable contraction of the earth's volume, and that we therefore have no just reason to reject the late determination of secular acceleration arising from a change of the eccentricity of the earth's orbit, because it does not cover the whole amount determined from the observations of ancient eclipses.

Professor Whitney read a paper on the progress of the Geological Survey of California, and presented to the Academy a volume of his report on the Paleontology of that State.

On the motion of Dr. H. R. Storer a committee was appointed, consisting of Professor Rogers, Dr. A. A. Gould, and Professor Agassiz, to consider what action the Academy should take toward promoting the Geological Survey of California and the adjacent territories.

---

**Five hundred and forty-fourth Meeting.**

January 10, 1865. — MONTHLY MEETING.

The PRESIDENT in the chair.

The Corresponding Secretary read letters relative to exchanges; also a communication from Mr. Otto Struve of the Observatory at Pulkova, announcing the decease of the eminent astronomer his father.

The following communication from Mr. E. W. Morley to Professor A. Hopkins of Williams College, was presented.

I have lately completed the reduction of my observations for the latitude of Williams College Observatory, and thinking you might like to have the result in some form which you can preserve, I transmit the final equations of condition with their solution. You will find the mode of observation fully described in Loomis's Practical Astronomy; briefly it is as follows. The telescope of the transit instrument being adjusted so as to move nearly in the plane of the prime vertical, the star's time of transiting the prime vertical is computed, together with its altitude at that time, and the transit pointed in readiness. The star moves transversely across the field, and is kept in the centre by moving the eyepiece horizontally and the telescope vertically. Its transits over the first ten wires are noted by the chronograph, the instrument is carefully reversed and the transits over the same wires in a reverse order are noted. After passing the meridian the transits over the same wires are noted again, and the instrument reversed as before. This gives forty passages to be noted, and, as a rule, ninety-six readings of the level. As the interval between the east and west passages is from three to five hours, there is but a small probability of getting a complete observation, and the loss of one of the four transits over each wire makes an observation valueless. Unfortunately, during the time I was occupied with the enclosed observations, the weather was so variable that a great proportion of my observations were lost, seven eighths or more I should think. The form of reduction is as follows. Take an observation of  $\alpha$  Lyrae, July 19, 1861. Transit over wire (1).

|                                    | h. | m. | s.               |
|------------------------------------|----|----|------------------|
| East vertical, circle South . . .  | 16 | 29 | 3.31             |
| “ “ North . . .                    | 16 | 35 | 2.51             |
| West vertical, circle North . . .  | 20 | 28 | 26.00            |
| “ “ South . . .                    | 20 | 34 | 23.57            |
| W. — E., circle South . . .        | 4  | 5  | 20.26            |
| W. — E., circle North . . .        | 3  | 53 | 23.49            |
| $\frac{1}{4}$ Sum . . .            | 1  | 59 | 40.94            |
| $\frac{1}{4}$ Difference . . .     | 2  | 59 | 19               |
| Cos $\frac{1}{4}$ Sum . . .        |    |    | 9.9378777        |
| Cos $\frac{1}{4}$ Difference . . . |    |    | 9.9999632        |
| Cotang $d$ (declination) . . .     |    |    | <u>0.0969342</u> |
| Cotang $\lambda$ (latitude) . . .  |    |    | 0.0347751        |

$$\begin{aligned} \lambda &= 42^{\circ} 42' 30''.73 \\ \text{Correction for level} &+3.30 \\ \text{" for azimuth} &-0.35 \\ \text{" for declination} &+1.02 \\ \lambda &42 42 34.75 \end{aligned}$$

The great part of my observations were lost by clouds, which would come up unexpectedly and spoil the labor of four hours, again and again. I have on some nights made as many as twenty observations, involving four hundred passages in the E. V., without getting a single star in the W. V., thus losing all. I can, therefore, rest the latitude of Williams College on only fifty-two observations, but hope these may be enough for present purposes.

Let  $\lambda' = (\text{latitude of P. V. instrument} - 42^{\circ} 42' 20'')$ .

$x$  be the value of one division of the level, which Mr. Phelps thinks to be near  $1''.0$ .

Values of  $a\lambda' + a'x - b = 0$ .

(A.) (*From Observation.*)

|    |                              | $\Delta$ | $\Delta^2$ |
|----|------------------------------|----------|------------|
| 1  | $\lambda' + 5.39x - 20.88 =$ | $- .92$  | .8464      |
| 2  | $\lambda' + 5.39x - 19.53 =$ | $- 2.27$ | 5.1529     |
| 3  | $\lambda' + 5.39x - 19.53 =$ | $- 2.27$ | 5.1529     |
| 4  | $\lambda' + 5.39x - 20.73 =$ | $- 1.07$ | 1.1449     |
| 5  | $\lambda' + 5.39x - 21.56 =$ | $- .24$  | .0576      |
| 6  | $\lambda' + 5.39x - 21.09 =$ | $- .71$  | .5041      |
| 7  | $\lambda' + 5.39x - 21.85 =$ | $+ .05$  | .0025      |
| 8  | $\lambda' + 5.39x - 22.27 =$ | $+ .47$  | .2209      |
| 9  | $\lambda' + 5.39x - 23.63 =$ | $+ 1.83$ | 3.3469     |
| 10 | $\lambda' + 5.39x - 21.64 =$ | $- .16$  | .0256      |
| 11 | $\lambda' + 5.38x - 21.85 =$ | $+ .06$  | .0036      |
| 12 | $\lambda' + 5.38x - 22.55 =$ | $+ .76$  | .5776      |
| 13 | $\lambda' + 5.38x - 20.73 =$ | $- 1.06$ | 1.1236     |
| 14 | $\lambda' + 5.38x - 21.40 =$ | $- .39$  | .1521      |
| 15 | $\lambda' + 5.38x - 21.04 =$ | $- .75$  | .5625      |
| 16 | $\lambda' + 5.38x - 22.06 =$ | $+ .27$  | .0727      |
| 17 | $\lambda' + 5.38x - 22.72 =$ | $+ .95$  | .9025      |
| 18 | $\lambda' + 5.38x - 23.02 =$ | $+ 1.23$ | 1.5129     |
| 19 | $\lambda' + 5.38x - 22.63 =$ | $+ .84$  | .7056      |
| 20 | $\lambda' + 6.40x - 23.35 =$ | $+ .57$  | .3249      |
| 21 | $\lambda' + 6.40x - 23.16 =$ | $+ .38$  | .1444      |
| 22 | $\lambda' + 6.40x - 22.07 =$ | $- .71$  | .5041      |

|   |                               | $\Delta$ | $\Delta^2$ |
|---|-------------------------------|----------|------------|
| 23  | $\lambda' + 6.40 x - 22.92 =$ | $+ .14$  | .0196      |
| 24  | $\lambda' + 6.40 x - 23.72 =$ | $+ .94$  | .8836      |
| 25  | $\lambda' + 6.40 x - 23.21 =$ | $+ .43$  | .1849      |
| 26  | $\lambda' + 6.40 x - 22.52 =$ | $- .26$  | .0676      |
| 27  | $\lambda' + 6.40 x - 24.87 =$ | $+ 2.09$ | 4.3681     |
| 28  | $\lambda' + 6.40 x - 23.32 =$ | $+ .54$  | .2916      |
| 29  | $\lambda' + 6.40 x - 23.22 =$ | $+ .44$  | .1936      |
| 30  | $\lambda' - 3.30 x - 10.40 =$ | $- 1.95$ | 8.8025     |
| 31  | $\lambda' - 3.30 x - 11.09 =$ | $- 2.26$ | 5.1136     |
| 32  | $\lambda' - 3.30 x - 11.26 =$ | $- 2.09$ | 4.3681     |
| 33  | $\lambda' - 3.30 x - 10.25 =$ | $- 3.10$ | 9.6100     |
| 34  | $\lambda' - 3.30 x - 11.42 =$ | $- 1.93$ | 3.7249     |
| 35  | $\lambda' - 3.30 x - 10.95 =$ | $- 2.40$ | 5.7600     |
| 36  | $\lambda' - 3.30 x - 10.62 =$ | $- 2.73$ | 7.4529     |
| 37  | $\lambda' - 3.30 x - 11.12 =$ | $- 2.23$ | 4.9729     |
| 38  | $\lambda' - 3.30 x - 9.75 =$  | $- 3.60$ | 12.9600    |
| 39  | $\lambda' - 3.30 x - 11.69 =$ | $- 2.66$ | 7.0756     |
| 40  | $\lambda' - 3.69 x - 14.26 =$ | $+ 1.29$ | 1.6641     |
| 41  | $\lambda' - 3.69 x - 14.37 =$ | $+ 1.40$ | 1.9600     |
| 42  | $\lambda' - 3.69 x - 15.72 =$ | $+ 2.75$ | 7.5625     |
| 43  | $\lambda' - 3.69 x - 15.93 =$ | $+ 2.96$ | 8.7816     |
| 44  | $\lambda' - 3.69 x - 15.82 =$ | $+ 2.85$ | 8.1225     |
| 45  | $\lambda' - 3.69 x - 15.34 =$ | $+ 2.37$ | 5.6169     |
| 46  | $\lambda' - 3.69 x - 15.20 =$ | $+ 2.23$ | 4.9729     |
| 47  | $\lambda' - 3.69 x - 15.26 =$ | $+ 2.29$ | 5.2441     |
| 48  | $\lambda' - 3.80 x - 13.40 =$ | $+ .53$  | .2809      |
| 49  | $\lambda' - 3.80 x - 15.15 =$ | $+ 2.28$ | 5.1984     |
| 50  | $\lambda' - 3.80 x - 13.76 =$ | $+ .89$  | .7921      |
| 51  | $\lambda' - 3.80 x - 13.95 =$ | $+ 1.06$ | 1.1236     |
| 52  | $\lambda' - 3.80 x - 13.33 =$ | $+ .46$  | .2116      |
| <b>(a.)</b> $52 \lambda' + 84.10 x - 944''.11 = 0, \quad \Sigma \Delta^2 = 145.4221.$ |                               |          |            |

Values of  $a' a \lambda' + a'^2 x - a' b = 0.$

| <b>(B.)</b> |                 |               |        |       |
|-------------|-----------------|---------------|--------|-------|
| 1           | 5.39 $\lambda'$ | $+ 29.05 x -$ | 112.54 | $= 0$ |
| 2           | 5.39 $\lambda'$ | $+ 29.05 x -$ | 105.27 | $= 0$ |
| 3           | 5.39 $\lambda'$ | $+ 29.05 x -$ | 105.27 | $= 0$ |
| 4           | 5.39 $\lambda'$ | $+ 29.05 x -$ | 111.73 | $= 0$ |
| 5           | 5.39 $\lambda'$ | $+ 29.05 x -$ | 116.21 | $= 0$ |
| 6           | 5.39 $\lambda'$ | $+ 29.05 x -$ | 113.68 | $= 0$ |
| 7           | 5.39 $\lambda'$ | $+ 29.05 x -$ | 117.77 | $= 0$ |
| 8           | 5.39 $\lambda'$ | $+ 29.05 x -$ | 120.03 | $= 0$ |

|    |   |      |    |   |       |   |   |        |    |   |
|----|---|------|----|---|-------|---|---|--------|----|---|
| 9  | + | 5.39 | λ' | + | 29.05 | x | — | 130.37 | == | 0 |
| 10 | + | 5.39 | λ' | + | 29.05 | x | — | 116.64 | == | 0 |
| 11 | + | 5.38 | λ' | + | 29.05 | x | — | 117.55 | == | 0 |
| 12 | + | 5.38 | λ' | + | 28.94 | x | — | 121.32 | == | 0 |
| 13 | + | 5.38 | λ' | + | 28.94 | x | — | 111.53 | == | 0 |
| 14 | + | 5.38 | λ' | + | 28.94 | x | — | 115.13 | == | 0 |
| 15 | + | 5.38 | λ' | + | 28.94 | x | — | 113.39 | == | 0 |
| 16 | + | 5.38 | λ' | + | 28.94 | x | — | 118.68 | == | 0 |
| 17 | + | 5.38 | λ' | + | 28.94 | x | — | 122.23 | == | 0 |
| 18 | + | 5.38 | λ' | + | 28.94 | x | — | 123.85 | == | 0 |
| 19 | + | 5.38 | λ' | + | 28.94 | x | — | 121.75 | == | 0 |
| 20 | + | 6.40 | λ' | + | 40.96 | x | — | 149.44 | == | 0 |
| 21 | + | 6.40 | λ' | + | 40.96 | x | — | 148.42 | == | 0 |
| 22 | + | 6.40 | λ' | + | 40.96 | x | — | 141.25 | == | 0 |
| 23 | + | 6.40 | λ' | + | 40.96 | x | — | 146.69 | == | 0 |
| 24 | + | 6.40 | λ' | + | 40.96 | x | — | 151.81 | == | 0 |
| 25 | + | 6.40 | λ' | + | 40.96 | x | — | 148.54 | == | 0 |
| 26 | + | 6.40 | λ' | + | 40.96 | x | — | 144.13 | == | 0 |
| 27 | + | 6.40 | λ' | + | 40.96 | x | — | 159.17 | == | 0 |
| 28 | + | 6.40 | λ' | + | 40.96 | x | — | 149.25 | == | 0 |
| 29 | + | 6.40 | λ' | + | 40.96 | x | — | 148.61 | == | 0 |
| 30 | — | 3.30 | λ' | + | 10.89 | x | + | 37.62  | == | 0 |
| 31 | — | 3.30 | λ' | + | 10.89 | x | + | 36.60  | == | 0 |
| 32 | — | 3.30 | λ' | + | 10.89 | x | + | 37.16  | == | 0 |
| 33 | — | 3.30 | λ' | + | 10.89 | x | + | 33.83  | == | 0 |
| 34 | — | 3.30 | λ' | + | 10.89 | x | + | 37.69  | == | 0 |
| 35 | — | 3.30 | λ' | + | 10.89 | x | + | 36.14  | == | 0 |
| 36 | — | 3.30 | λ' | + | 10.89 | x | + | 35.05  | == | 0 |
| 37 | — | 3.30 | λ' | + | 10.89 | x | + | 36.70  | == | 0 |
| 38 | — | 3.30 | λ' | + | 10.89 | x | + | 32.18  | == | 0 |
| 39 | — | 3.30 | λ' | + | 10.89 | x | + | 38.57  | == | 0 |
| 40 | — | 3.69 | λ' | + | 13.62 | x | + | 52.62  | == | 0 |
| 41 | — | 3.69 | λ' | + | 13.62 | x | + | 54.03  | == | 0 |
| 42 | — | 3.69 | λ' | + | 13.62 | x | + | 58.01  | == | 0 |
| 43 | — | 3.69 | λ' | + | 13.62 | x | + | 58.78  | == | 0 |
| 44 | — | 3.69 | λ' | + | 13.62 | x | + | 58.38  | == | 0 |
| 45 | — | 3.69 | λ' | + | 13.62 | x | + | 56.60  | == | 0 |
| 46 | — | 3.69 | λ' | + | 13.62 | x | + | 56.09  | == | 0 |
| 47 | — | 3.69 | λ' | + | 13.62 | x | + | 56.21  | == | 0 |
| 48 | — | 3.80 | λ' | + | 14.44 | x | + | 50.92  | == | 0 |
| 49 | — | 3.80 | λ' | + | 14.44 | x | + | 57.57  | == | 0 |
| 50 | — | 3.80 | λ' | + | 14.44 | x | + | 52.29  | == | 0 |
| 51 | — | 3.80 | λ' | + | 14.44 | x | + | 53.01  | == | 0 |
| 52 | — | 3.80 | λ' | + | 14.44 | x | + | 50.65  | == | 0 |

(b.)  $85.10 \lambda' + 1250.65 x - 2625''.55 = 0.$

|         |                     |               |               |                     |
|---------|---------------------|---------------|---------------|---------------------|
| (a)     | 52.00 $\lambda'$ +  | 85.10 $x$ -   | 944".11 = 0,  |                     |
| (b)     | 85.10 $\lambda'$ +  | 1250.65 $x$ - | 2625".55 = 0, |                     |
| log     | 1250.65             | 3.0971358     |               | - 13874.87          |
| log     | 85.10               | 1.9299296     |               | + 2625.55           |
| log (a) |                     | 1.1672062     |               | - 11249.32          |
| log     | 52                  | 1.7160035     |               | 1250.65 $x$         |
| log     | 52 + log (a)        | 2.8832095     | 764.20        | - 1250.65 $x$       |
|         |                     |               |               | 0.0 $x$             |
| log     | 944.11              | 2.9750226     |               |                     |
| log     | (944.11 $\times$ a) | 4.1422288     | 13874.87      | + 764.20 $\lambda'$ |
|         |                     |               |               | - 85.10 $\lambda'$  |

Hence,  $764''.20 \lambda' + 1250''.65 x - 13874''.87 = 0 + 679''.10 \lambda'$   
 $679.10 \lambda' - 11249.32 = 0.$

|     |            |           |                      |     |
|-----|------------|-----------|----------------------|-----|
| log | 11249".32  | 4.0511263 |                      |     |
| log | 679".10    | 2.8319337 |                      |     |
| log | $\lambda'$ | 1.2191926 | $\lambda' = 16''.56$ | (1) |

|     |                      |           |                |            |
|-----|----------------------|-----------|----------------|------------|
| log | 85".10               | 1.9299296 |                |            |
| log | (85".10 $\lambda'$ ) | 3.1491222 | 1409".80       | - 1409".80 |
|     |                      |           |                | 2625".55   |
| log | 1215".75             | 3.0848452 |                | 1215".75   |
| log | 1250".65 ( $x$ )     | 3.0971358 |                |            |
| log | ( $x$ )              | 9.9877064 | $x = 0''.9721$ | (2)        |

|                               |                          |                                      |                                |
|-------------------------------|--------------------------|--------------------------------------|--------------------------------|
| $\Sigma \Delta^2 = 145.4221,$ |                          | $\frac{\Sigma \Delta^2}{52} = 2.603$ |                                |
| log                           | 2.603                    | 0.4156910                            |                                |
| log                           | (2.603) $^{\frac{1}{2}}$ | 0.2078455                            |                                |
| log                           | constant                 | 9.8289751                            | Probable error of single       |
| log                           | prob. error              | 0.0368206                            | observation $\pm 1''.0885$ (3) |

|     |                          |         |                        |
|-----|--------------------------|---------|------------------------|
| log | (2.603) $^{\frac{1}{2}}$ | 0.20785 |                        |
| log | constant                 | 9.60091 |                        |
| log | mean error               | 9.80876 | Mean error 0''.644 (4) |

|     |  |         |                     |
|-----|--|---------|---------------------|
| log | $\left(\frac{2 \Sigma \Delta^2}{n}\right)$ | 0.71650 |                     |
| log | reciprocal                                 | 9.28350 | Precision .1921 (5) |
| log | $\sqrt{\text{reciprocal}}$                 | 9.64175 | Weight .4383 (6)    |

|                              |  |                                  |
|------------------------------|--|----------------------------------|
|                              | $\frac{\Sigma \Delta^2}{52-2} = e^2 = 2.908442$                  |                                  |
| $\log e^2$                   | 0.463660   |                                  |
| $\log \chi^2$                | 0.825686   | Table in Coast Survey Report,    |
| $\log \chi^2 e^2$            | 1.289346   | 1854, p. 136.*                   |
| $\log \chi e$                | .644673  |                                  |
| $\chi e$                     | $\pm 4''.412$  | Limit by Peirce's Criterion. (7) |
| $\Sigma a^2 = 52$            | $\Sigma a'^2 = 1250.65$  | $\Sigma a a' = 85.10$            |
| $(\Sigma a a')^2 = 7242.01$  | $c = \Sigma a^2 \cdot \Sigma a'^2 - (\Sigma a a')^2 = 57791.79.$ |                                  |
|                              | $\lambda'.$  | $x.$                             |
| $\log \Sigma a'^2$           | 3.096118   | $\log \Sigma a^2$ 1.71600        |
| $\log c$                     | 4.761868   | 4.76187                          |
| $\log d^2$                   | 8.334250   | $\log d'^2$ 6.95413              |
| $\log d$                     | 9.167125   | $\log d'$ 8.47706                |
| $\log$ mean error            | 9.80876  | 9.80876                          |
| $\log$ mean error of result  | 8.97589  | 8.28582                          |
|                              | Mean error $\lambda' 0''.0946$ (8)                               | Mean error $x 0''.0193$ (9)      |
| $\log$ constant              | 0.22807  | 0.22807                          |
| $\log$ prob. error of result | 9.20396  | 8.51389                          |
|                              | Probable error $\lambda' 0''.1599$ (10)                          | Prob. error $x 0''.0327$ (11)    |

(A) are the fifty-two observations. Each equation should be  $= 0$ , the quantities in the column  $\Delta$  are the errors of each observation. The column  $\Delta^2$  contains the squares of these errors, and the values of  $\lambda'$  and  $x$  are those which make  $\Sigma \Delta^2$  (the sum of the squares of the errors) a minimum, and are, therefore, the *most probable* values. (3) gives the quantity  $1''.0885$ , as the probable error of a single observation. (7) gives  $4''.41$  as the limit, by Peirce's Criterion, for a doubtful observation. (10) gives, as the probable error of the resulting latitude,  $0''.16$ . It is more likely than not that the result is within this amount of the truth. The common tables will show that it is 100 to 1 that my latitude is within  $0''.61$ ; 200 to 1 that it is within  $0''.67$ ; and 300 to 1 that it is within  $0''.78$  of the truth.

The prime vertical transit was one hundred feet south of the centre of the dome, which is  $1''.0$  of arc, which is to be added to my result.  $\alpha$  Lyræ's place was taken from the English Nautical Almanac. Its declination ( $d$ ) is  $1''.0$  different in the American, and  $0''.1$  less in the



Berlin Jahrbuch. To reduce to either of these authorities, the variation of the declination ( $\delta d$ ) must be multiplied into 1.02. When all terms are taken into account, the latitude is

$$\lambda = 42^\circ 42' 20'' + 16''.56 + 1''.0 + 1.02 \delta d \pm 0''.1599,$$

from which for use,

$$\lambda = 42^\circ 42' 37''.56 \quad \text{or,} \quad 42^\circ : 42' : 37''.6 :$$

The value of one division of the level is  $0''.972 \pm ''0.033$ .

**Five hundred and forty-fifth Meeting.**

January 25, 1865.—STATUTE MEETING.

The PRESIDENT in the chair.

The President called the attention of the Academy to the recent decease of two of its members,—Hon. Edward Everett and Mr. W. P. G. Bartlett, of the Resident Fellows.

Professor Henry D. Rogers, of Glasgow, was elected an Associate Fellow in Class II. Section 1.

Mr. Frederick W. Putnam, of Salem, was elected a Resident Fellow in Class II. Section 3.

**Five hundred and forty-sixth Meeting.**

February 14, 1865.—MONTHLY MEETING.

The PRESIDENT in the chair.

The President called the attention of the Academy to the recent decease of Capt. J. M. Gilliss, an Associate Fellow, and the Director of the National Observatory at Washington.

Mr. Ferrel made the following communication:—

“It was shown in a note read on a former occasion, that if the tidal wave of the ocean is on an average two feet high and displaced two degrees by friction, the effect of the moon’s attraction on the tidal wave must cause an increase of  $\frac{1}{140}$  of a second in the length of the day in 2,500 years. From this we may readily determine that the earth’s